DSC 140A - Homework 04

Due: Wednesday, February 7

Write your solutions to the following problems by either typing them up or handwriting them on another piece of paper. Unless otherwise noted by the problem's instructions, show your work or provide some justification for your answer. Homeworks are due via Gradescope at 11:59 PM.

Problem 1.

In this problem, you will derive the solution to the ridge regression optimization problem.

Recall that the ridge regression regularized risk function is

$$\tilde{R}(\vec{w}) = \frac{1}{n}\sum_{i=1}^{n}(\vec{\phi}(\vec{x}^{(i)}) \cdot \vec{w} - y_i)^2 + \lambda\|\vec{w}\|^2.$$

Here, $\vec{\phi}(\vec{x})$ is a feature map. Also recall that this risk function can be equivalently written in matrix-vector form as

$$\tilde{R}(\vec{w}) = \frac{1}{n}\|\Phi\vec{w} - \vec{y}\|^2 + \lambda\|\vec{w}\|^2,$$

where Φ is the design matrix; its ith row is $\vec{\phi}(\vec{x}^{(i)})$.

a) Show that $\tilde{R}(\vec{w}) = \frac{1}{n}\left(\vec{w}^T\Phi^T\Phi\vec{w} - 2\vec{w}^T\Phi^T\vec{y} + \vec{y}^T\vec{y}\right) + \lambda\vec{w}^T\vec{w}$.

b) So far this quarter, we have seen a few vector calculus identities. For example, we know that $\frac{d}{d\vec{w}}(\vec{w}^T\vec{w}) = 2\vec{w}$.

Using these identities, show that

$$\frac{d}{d\vec{w}}\tilde{R}(\vec{w}) = \frac{1}{n}\left(2\Phi^T\Phi\vec{w} - 2\Phi^T\vec{y}\right) + 2\lambda\vec{w}.$$


c) Show that the minimizer of $\tilde{R}(\vec{w})$ is $\vec{w}^* = (\Phi^T\Phi + n\lambda I)^{-1}\Phi^T\vec{y}$.

Problem 2.

The data set linked below contains data for performing non-linear regression. The first column is x (the independent variable), and the second column is y (the dependent variable).

`https://f000.backblazeb2.com/file/jeldridge-data/010-nonlinear_regression/data.csv`

Plotting the data shows that there is a non-linear relationship between x and y:



a) Fit a function of the form $H(\vec{x}) = w_1\phi_1(\vec{x}) + w_2\phi_2(\vec{x}) + \ldots + w_{50}\phi_{50}(\vec{x})$, where each $\phi_i(\vec{x})$ is a Gaussian basis function. Your 50 Gaussian basis functions should be equally-spaced, with the first at $\mu_1 = -10$ and the last at $\mu_{50} = 10$. The width of each Gaussian should be $\sigma = 1$. You should *not* augment $\vec{x}$.

For your answer, report only w_1 (the first component of $\vec{w}$), and show your code.

b) Plot the prediction function $H(\vec{x})$ that you trained in the previous part, on top of the data. Provide your plot and the code used to generate it.

c) You should see that the prediction function $H(\vec{x})$ slighly overfits the data. Now perform ridge regression on the same data, using the same Gaussian basis functions. Choose the regularization parameter λ to reduce overfitting (you may do so by trial and error – no need to perform cross-validation). For your answer, state λ and plot your new prediction function on top of the data. Also provide your code.